



TrueHume™ Fulvic70 0-0-3

Features and benefits

TrueHume Fulvic70 0-0-3 is 100% water-soluble and can be used as a soil or foliar application. It may enhance nutrient uptake by plants. It is compatible with most powdered and liquid fertilizers, biological and enzymatic formulations. can be blended with granular fertilizer or dissolved in water to produce a concentrated liquid.

GUARANTEED MINIMUM ANALYSIS

Soluble Potash (K₂O) 3%
 Fulvic Acid 70%
 Derived From: Peat

APPLICATION RATES

VEGETABLES & FLOWERS, GREENHOUSE & FIELD	Foliar Application: 0.5-2 kg/ha (0.2-0.8 kg/ac) - Root/Drip Irrigation: 1-3 kg/ha (0.4-1.2 kg/ac)				
	First	Second	Third	Optional	
Melons, eggplants, peppers, squash	Six-leaf stage	Early bloom	First fruit set	Four weeks later	
Beans, peas, lentils, canola	Six-leaf stage	First bloom	First pods		
Broccoli, cabbage, cauliflower	Six true leaf stage	Three weeks later	Head initiation		
Cucumbers, cucurbits	Six-leaf stage	Just prior to bloom	Whilst picking		
Sweet corn	Six-leaf stage	55-75cm/20-30 inch growth	Just prior to tasseling		
Leeks, onions, carrots, turnips	2-3 weeks after emergence	Root enlargement	Every 2 weeks until harvest		
Potatoes	Six-leaf stage	Approx. Pea size	Early bloom	Bulking up	
Tomatoes (fresh fruit market)	Six-leaf stage	Early bloom	Whilst picking	After 2 weeks	
Leafy vegetables, lettuce, etc.	Every 2 weeks				
Nursery plants, containers & field	Every 2 weeks				
Hydroponic gardens and cannabis	Mix 1ml of fulvic acid concentrate (see below for mixing instructions) per liter of water in reservoir				
FRUIT	Foliar Application: 0.5-2 kg/ha (0.2-0.8 kg/ac) - Root/Drip Irrigation: 1-3 kg/ha (0.4-1.2 kg/ac)				
	First	Second	Third	Fourth	Optional
Grapes	20-30cm cane	45-60cm cane	Full blooming date	Berry set/early shattering	3 weeks later
Apples/pears	Green tip	Pre-bloom pink bud	Full blooming date	Early fruit formation	Every 3 weeks
Strawberries & other berries	Just after transplant	Prior to bloom	Whilst picking	3 weeks later	Every 3 weeks
Stone fruit	Bud burst	Petal fall	4 weeks later	3 weeks later	Every 3 weeks
Plums, cherries	Pink white bud	Full bloom	Early fruit formation	3 weeks later	Every 3 weeks
ECONOMIC CROP	Foliar Application: 0.5-2 kg/ha (0.2-0.8 kg/ac) - Root/Drip Irrigation: 1-3 kg/ha (0.4-1.2 kg/ac)				
	First	Second	Third	Optional	
Oilseed crops	Before reproductive growth stage	At height of flowering	Every 4 weeks		
Tobacco	Early post emergence	After every leaves picked	Early bloom	Every 4 weeks	
Hops	Early post emergence	Five weeks later	Every 4 weeks		
Forage crops	After each cut				
TURF	Foliar Application				
1-4 KG/HA	1. Regular applications to greens and tees are especially beneficial 2. Applied as bio-stimulant to increase the number and activity of micro-organisms. This will improve grass growth and help reduce soil borne disease symptoms in the turf. 3. Tiller will also be increased but without excessive top growth				
GRAINS	Foliar Application: 1-3 kg/ha (0.4-1.2 kg/ac) - Root/Drip Irrigation: 2-6 kg/ha (0.8-2.4 kg/ac)				
	First	Second	Optional		
Winter wheat and oats	First node	Flag leaf	After any environmental stress		
Winter (malting) barley	Early post emergence	After any environmental stress			
Spring wheat and oats	1-3 tillers	First node	Flag leaf and after any environmental stress		
Spring (malting) barley	1-3 tillers	First node	Flag leaf and after any environmental stress		
Corn	2-6 leaf stage	At 50-75 cm. Growth	Just prior to tasseling		

SEED TREATMENT: 0.5 kg per tonne of seed

Fulvic acid powder can be blended with granular fertilizer or dissolved in water to produce a concentrated liquid.

To make a 10% Fulvic Concentrate solution - Add 150 g/litre or 15 kg/100 litres of TrueHume Fulvic70 0-0-3 to water and mix vigorously.

WARNING

Keep out of reach of children and animals. Harmful if swallowed. Harmful in contact with skin. Harmful if inhaled. Wear protective equipment including clothing, gloves, and eye and respiratory protection when using this product. Do not eat, drink or smoke when using this product. Avoid breathing dust. Use only in a well-ventilated location. Wash thoroughly after handling. If exposed, concerned, or feeling unwell seek medical attention. If swallowed, rinse mouth. If on skin, wash thoroughly with water and wash clothing before reuse. If inhaled, move person to fresh air and keep comfortable. Store in original container and out of reach of children. Dispose of contents and container in accordance with local, regional, and national regulations.

25 KG

Lot Number:

Expiry Date:

Registration Number 2019131A Fertilizers Act

Packaged by Agriculture Solutions Inc. 2079 Line 34, Shakespeare ON, N0B 2P0

855 247 6548 | www.agsolcanada.com

